

中國螟蛾科昆蟲名錄

胡氏「中國昆蟲目錄」補遺

部分二. 水螟、苔螟及擬螟亞科

陸近仁

(北京農業大學, 中國科學院)

管致和

(北京農業大學)

水螟亞科 *Nymphulinae* Fernald 1902

Aediodina quaternalis (Lederer, 1863)

台灣 (Shibuya, 1938; Klima, 1937).

Ambia albobasalis Caradja & Meyrick, 1934

華東 (Klima, 1937); 廣東 (Caradja & Meyrick, 1934).

A. amoyalis Caradja, 1925

華南 (Klima, 1937); 福建、廣東 (Caradja, 1925).

A. colonalis (Bremer, 1864)

台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja & Meyrick, 1934; Klima, 1937).

A. likiangalis Caradja, 1937

雲南 (Caradja, 1937).

A. marginalis Moore, 1887

華東、華南 (Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

A. melanistis Hampson, 1917

台灣 (Shibuya, 1928; Klima, 1937).

Aulacodes laminalis South, 1901

中國 (South, 1901); 華中、華西 (Caradja, 1925; Klima, 1937); 江西、福建、廣東、廣州 (Caradja, 1925; Caradja & Meyrick, 1934).

Au. mimeticalis Caradja, 1925

華南 (Klima, 1937); 福建、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Au. ochripicta (Moore, 1887)

華南、華西 (Klima, 1937); 江西、福建、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Au. reductalis Caradja, 1925

華南 (Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Au. simplicialis f. *rufalis* Caradja, 1938

福建 (Caradja, 1938).

Au. sinensis Hampson, 1897

華西 (Caradja, 1925; Klima, 1937); 四川 (Hampson, 1897; Klima, 1937).

Bradina admixtal (Walker, 1859)

華南、華西 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 江蘇、浙江 (李鳳藻, 1952).

B. atopal (Walker, 1859)

華北、華西 (Klima, 1937); 華中 (Caradja, 1925; Klima, 1937); 華東、北京、浙江 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

B. concordalis (Lederer, 1863)

華南 (Klima, 1937); 台灣、廣州 (Caradja, 1925 (var.)); 廣東 (Caradja & Meyrick, 1934).

B. concordalis var. *defloralis* Strand, 1919

台灣 (Shibuya, 1928; Klima, 1937).

B. consortialis Caradja, 1927

四川 (Klima, 1937).

B. eriltoides Strand, 1919

台灣 (Shibuya, 1928; Klima, 1937).

B. geminalis Caradja, 1927

廣東 (Caradja & Meyrick, 1934; Klima, 1937); 四川 (Klima, 1937).

*B. intermedi*alis Caradja, 1932

華南 (Caradja, 1932).

B. megesalis (Walker, 1859)

華北 (Klima, 1937); 華中、華西 (Caradja, 1925; Klima, 1937); 華南、上海 (Caradja, 1925).

B. melanoperas Hampson, 1896

華西、華中、華南 (Klima, 1937); 浙江、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

B. melanoperas *sinensis* Caradja, 1925

華中、華西、華南 (Caradja, 1925); 浙江、廣東、廣州 (Klima, 1937).

B. nigripunctalis South, 1901

中國 (South, 1901); 華西 (Caradja, 1925; Klima, 1937); 廣東 (Caradja, 1925; Caradja, & Meyrick, 1934); 廣州 (Caradja, 1925).

B. rectilinealis South 1901

中國 (South, 1901); 華中、華西 (Klima, 1937).

Camptomastix hisbonalis (Walker, 1859)

中國 (Hampson, 1896); 華東 (Caradja, 1925); 華中 (Caradja, 1925; Klima, 1937); 華西 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja & Meyrick, 1934).

Cataclysta angulata Moore, 1885

台灣 (Shibuya, 1928; Klima, 1937).

C. blandialis Walker, 1859

華南 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

C. blandialis f. *nigritalis* Hampson, 1893

華東、華南 (Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

- C. croesusalis* Walker, 1859
廣東 (Caradja & Meyrick, 1934; Klima, 1937).
- C. croesusalis angustalis* Caradja, 1925
廣州 (Caradja, 1925; Klima, 1937).
- C. fuscalis* Hampson, 1893
廣東 (Klima, 1937); 華南 (Caradja, 1925).
- C. midas* Butler, 1881
華中等 (Caradja, 1925; Klima, 1937).
- Clupeosoma polusalis* (Walker, 1859)
華東等 (Klima, 1937); 廣東 (Caradja & Meyrick, 1934).
- Cymoriza hoenei* Caradja & Meyrick, 1934
廣東 (Caradja & Meyrick, 1934; Klima, 1937).
- Cy. interruptalis* (Wileman & South, 1917)
台灣 (Shibuya, 1928; Klima, 1937).
- Cy. irrectalis* Guenée, 1854
華南 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Cy. taiwanalis* Shibuya, 1928
台灣 (Shibuya, 1928; Klima, 1927).
- Daulia afralis* Walker, 1859
華東、華南 (Klima, 1937) 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Dectilogaster amoyalis* (Caradja, 1932)
華南 (Caradja, 1932); 福建 (Klima, 1937).
- D. angustalis* (Caradja & Meyrick, 1934)
廣東 (Caradja & Meyrick, 1934 (*Tatobotys*); Klima, 1937).
- D. depalpalis* (Strand, 1919)
台灣 (Shibuya, 1928; Klima, 1937).
- D. janapalis* (Walker, 1859)
台灣 (Hampson, 1896 (*Tatobotys*); Shibuya, 1928; Caradja & Meyrick, 1934 (*Tatobotys*); Klima, 1937); 廣東 (Caradja & Meyrick, 1934; Klima, 1937).
- Diathrausta brevifascialis* (Wileman, 1911)
台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja & Meyrick, 1934).
- Di. picata* Butler, 1889
華東 (Caradja, 1925; Klima, 1937); 華南、華西 (Klima, 1937); 廣東 (Caradja, 1925).
- Di. plumbealis* Warren, 1896
華東 (Caradja, 1925; Klima, 1937); 華南、華西 (Klima, 1937); 江西 (Caradja, 1925).
- Di. plumbealis* var. *talboti* (Caradja, 1927)
中國 (Caradja, 1927).
- Di. profundalis* (Lederer, 1863)
華南 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925; Caradja & Mey-

rick, 1934).

Diatraustodes fulvofusa Hampson, 1901

華中等 (Caradja, 1925; Klima, 1937).

Eoöphila conjunctalis (Wileman & South, 1917)

台灣 (Shibuya, 1928 (*Eoöphyla*); Klima, 1937)

E. gibbosalis (Guenée, 1854)

台灣等 (Shibuya, 1928 (*Eoöphyla*); Klima, 1937).

E. peribocalis (Walker, 1859)

中國 (Hampson, 1896 (*Aulacodes*)); 華東、華中 (Caradja, 1925; Klima, 1937); 華西、華南 (Klima, 1937); 江西、福建 (Caradja, 1925); 廣東 (Caradja, 1925 (*Au.*); Caradja & Meyrick, 1934 (*Au.*)).

Eugauria albidentata (Hampson, 1897)

台灣 (Shibuya, 1928; Klima, 1937).

Eurrhpara urticata (Linnaeus, 1761)

華中、華西 (Caradja, 1925; Klima, 1937); 華南 (Klima, 1937); 雲南 (Caradja & Meyrick, 1934).

Gargela renatusalis Walker, 1859

華東、華南 (Klima, 1937); 廣東 (Caradja, 1925 (*Mixophila*); Caradja & Meyrick, 1934 (*M.*)).

Hyalophaga pulchralis var. *reducta* Caradja, 1931

四川 (Klima, 1937).

Luma ornatalis (Leech, 1899)

華西 (Caradja, 1925; Klima, 1937); 華中、華南 (Klima, 1937); 浙江、廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 上海 (Caradja, 1925).

Mabra eryxalis (Walker, 1859)

華南 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Massepha absolutalis Walker, 1859

台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Klima, 1937; Caradja & Meyrick, 1934 (*absolutalis*)).

M. ambialis Hampson, 1903

華南 (Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

M. rectangularis Caradja & Meyrick, 1934

中國 (Klima, 1937); 廣東 (Caradja & Meyrick, 1934).

M. talboti Caradja, 1927

華西 (Klima, 1937).

Musotima franckti Caradja, 1927

華西 (Klima, 1937).

M. nubilalis South, 1901

中國 (South, 1901); 華中 (Caradja, 1925; Klima, 1937).

M. suffusalis Hampson, 1893

華南 (Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Nymphula crisonalis (Walker, 1859)

華中 (Caradja, 1925); 華東、華南 (Klima, 1937); 浙江 (Hampson, 1896); 廣東 (Caradja, 1925;

Caradja & Meyrick, 1934).

N. defintialis (Strand, 1919)

台灣 (Shibuya, 1928; Klima, 1937).

N. depunctalis (Guenée, 1884)

華南 (Klima, 1937); 福建 (Caradja, 1925; 李鳳蓀, 1952); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1937; 李鳳蓀, 1952); 廣東 (Caradja & Meyrick, 1934).

N. diffualis nigrialbalis Caradja, 1925

廣東 (Caradja, 1925).

N. diminutalis (Snellen, 1880)

中國 (Hampson, 1896); 台灣 (Shibuya, 1928; Klima, 1928).

N. enixalis (Swinhoe, 1885)

華東 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja & Meyrick, 1934 (*erixalis*)).

N. fengshanalis (Pryer, 1877)

華北 (Klima, 1937); 華中 (Caradja, 1925; Klima, 1937); 華東 (Caradja & Meyrick, 1934; Klima, 1937); 浙江 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

N. fluctuosalis Zeller, 1852

華西 (Klima, 1937); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1937; Hampson, 1896); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣西 (李鳳蓀, 1952).

N. foederalis (Guenée, 1854)

華東 (Klima, 1937); 山東 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934 (*phoederalis*)).

N. foederalis f. cantonalis Caradja, 1925

廣州 (Caradja, 1925; Klima, 1937).

N. fulguralis Caradja, & Meyrick, 1934

廣東 (Caradja & Meyrick, 1934; Klima, 1937); 雲南、四川 (Caradja & Meyrick, 1934).

N. hampsoni (South, 1901)

華西 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937).

N. incurvalis South, 1901

中國 (South, 1901); 華中 (Caradja, 1925; Klima, 1937).

N. interruptalis (Pryer, 1877)

華東、華中、華南、華西 (Klima, 1937); 上海 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

N. interruptalis separatalis (Leech, 1889)

廣東 (Caradja & Meyrick, 1934 (var.)); 李鳳蓀, 1952); 江蘇、廣西 (李鳳蓀, 1952).

N. nigra Warren, 1896

台灣等 (Shibuya, 1928; Klima, 1937).

N. nigrolinealis Pryer, 1877

華中、華西 (Klima, 1937); 廣東 (Caradja, 1925 (f.)); Caradja & Meyrick, 1934 (f.)).

N. nymphaeala (Linnaeus, 1758)

華東 (Klima, 1937); 上海 (Caradja, 1925); 廣東、雲南 (Caradja & Meyrick, 1934).

- N. nymphaeata benesgnata* Caradja, 1925
華東、華中 (Klima, 1937); 湖北 (Caradja, 1925).
- N. sinicalis* Hampson, 1897
中國 (Hampson, 1897); 華東 (Caradja, 1925); 浙江 (Klima, 1937).
- N. turbata* (Butler, 1881)
華中 (Caradja, 1925; Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja & Meyrick, 1934 (*Parthenodes sutschana* Hampson)); 上海、福建、廣州 (Caradja, 1925 (*Parthenodes sutschana* Ragonot 1894)).
- N. ussuriensis* Rebel, 1910
華東、華南 (Klima, 1937); 上海、廣東 (Caradja & Meyrick, 1934); 廣西 (李鳳蓀, 1952).
- N. uxortalis* Strard, 1919
台灣 (Shibuya, 1928; Klima, 1937).
- N. vittalis* (Bremer, 1864)
華東、華中 (Caradja, 1925; Klima, 1937); 東北 (Klima, 1937); 廣東 (Caradja & Meyrick, 1934).
- Oligostigma bifurcalis* Pryer, 1877
華南 (Klima, 1937); 浙江 (Hampson, 1896 (*bifurcale*); Caradja, 1925 (*bifurcale*); Caradja & Meyrick, 1934 (*bifurcale*)); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- O. bifurcalis* (*bifurcale*) var. *szechuanalis* Caradja, 1935
四川 (Caradja, 1935).
- O. hapilista* (Swinhoe, 1892)
台灣等 (Shibuya, 1928; Klima, 1937).
- O. insectalis* Pryer, 1877
上海 (Klima, 1937; Caradja, 1925); (*insectale*)).
- O. kwangtungialis* Caradja, 1925
華南 (Klima, 1937); 江西、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- O. minutale* Caradja, 1932
華南 (Caradja, 1932).
- O. minutalis* Caradja, 1932
華南 (Klima, 1937).
- O. villidalis* Walker, 1859
台灣等 (Shibuya, 1928; Klima, 1937).
- Parthenodes aurantialis* Swinhoe, 1895
華東、華南 (Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- P. distinctalis* (Leech, 1889)
華中、華西 (Caradja, 1925; Klima, 1937); 廣東 (Caradja & Meyrick, 1934).
- P. distinctalis* var. *taiwanalis* (Wileman & South, 1917)
台灣 (Shibuya, 1928; Klima, 1937).
- P. inextricata* Moore, 1887
廣東 (Caradja & Meyrick, 1934; Klima, 1937).

- P. pallidalis* South, 1901
中國 (South, 1901); 華中 (Caradja, 1925; Klima, 1937).
- P. prodigalis* (Leech, 1889)
華中、華西 (Caradja, 1925; Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja & Meyrick, 1934).
- P. prodigalis* var. *cataclystalis* (Strand, 1919)
台灣 (Shibuya, 1928; Klima, 1937).
- P. prodigalis yunnanensis* Caradja, 1937
雲南 (Caradja, 1937).
- P. stellata* Warren, 1897
華中 (Caradja, 1925; Klima, 1937); 華南 (Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- P. triangularis* South, 1901
中國 (South, 1901); 華西 (Klima, 1937; Caradja, 1925); 浙江、雲南 (Caradja & Meyrick, 1934); 廣東 (Klima, 1937; Caradja & Meyrick, 1934).
- P. vagalis* (Walker, 1865)
廣東 (Caradja & Meyrick, 1934; Klima, 1937).
- Pelena bimaculalis* Shibuya, 1928
台灣 (Shibuya, 1928; Klima, 1937).
- Pe. obscuralis* (Swinhoe, 1895)
華中、華南 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937).
- Pe. phottas* (Meyrick, 1894)
台灣等 (Shibuya, 1928; Klima, 1937).
- Pe. sericea* (Bulter, 1869)
華中 (Caradja, 1925 (Luma); Klima, 1937); 華南 (Klima, 1937); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1937); 廣東 (Caradja & Meyrick, 1934 (Luma)).
- Perinephela doerriesti* Staudinger, 1892
華西 (Klima, 1937).
- Per. lancealis* (Schiffermüller & Denis, 1775)
華中 (Caradja, 1925); 華南、華西 (Klima, 1937); 廣東 (Caradja & Meyrick, 1934).
- Pileotocera aegimiusalis* (Walker, 1859)
華南 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937; Caradja, 1925 (f. *pallatalis*)); 福建 (Caradja, 1925).
- Pt. chrysorycta* Meyrick
華西 (Caradja, 1925).
- Pt. confinalis* (Caradja, 1927)
華西 (Klima, 1937).
- Pt. contingens* Moore
廣東 (Caradja & Meyrick, 1934).
- Pt. elongalis* (Warren, 1896)
台灣 (Shibuya, 1928; Klima, 1937).

Pi. immundalis (South, 1901)

湖北 (Klima, 1937).

Pi. sodalis (Leech, 1889)

華東等 (Caradja, 1925; Klima, 1937).

Psammotis hyalinalis Hübner, 1796

華中、阿爾泰山 (Klima, 1937); 上海等 (Caradja, 1925).

Ps. spissalis (Guenée, 1854)

台灣等 (Shibuya, 1928; Klima, 1937).

Stegothyris diagonalis (Guenée, 1854)

華南 (Klima, 1937); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925 (*Stenothyris*); Caradja & Meyrick, 1934).

Stenia amoenialis (Christoph, 1881)

華東、華南 (Klima, 1937); 浙江 (Caradja & Meyrick, 1934); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

S. charontalis (Walker, 1859)

華中、華西 (Klima, 1937; Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1937); 上海 (Caradja, 1925 (*Mabra*)); 廣東 (Caradja, 1925 (*Mabra*); Caradja & Meyrick, 1934).

S. minoralis (Snellen, 1880)

華南 (Klima, 1937); 台灣 (Caradja, 1925 (var.); Shibuya, 1928 (*spodinopa*); Caradja & Meyrick, 1934 (var.); Klima, 1937); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

S. nigriscripta (Swinhoe, 1895)

華南 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925 (*Mabra*); Caradja & Meyrick, 1934).

Strepsinoma hapilistalis (Strand, 1919)

台灣 (Shibuya, 1928; Klima, 1937).

Symphonia trivittalis (Warren, 1895)

台灣 (Shibuya, 1928; Klima, 1937).

Talanga sexpunctalis (Moore, 1877)

台灣 (Hampson, 1896; Caradja, 1925; Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Thysanoidma octalis Hampson, 1891

華南 (Klima, 1937); 台灣 (Shibuya, 1928; Klima, 1937); 廣東 (Caradja, 1925).

苔螟亞科 *Scopariinae* Ragonot 1890

Micraglossa flavidalis Hampson, 1907

華西 (Caradja, 1925; Klima, 1937).

M. scoparialis Warren, 1891

華西 (Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Scoparia ambigua (Treitschke, 1829)

華南 (Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

S. congestalis Walker, 1859

- 華中、華西 (Caradja, 1925; Klima, 1937); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
S. crataegella (Hübner, 1796)
 華西 (Klima, 1937).
S. frequentella Stainton, 1859
 華西 (Klima, 1937).
S. metaleucalis Hampson, 1907
 華西 (Klima, 1937).
S. murificalis Walker, 1859
 華西 (Caradja, 1925; Klima, 1937).
S. promiscua Wileman & South, 1919
 台灣 (Shibuya, 1928; Klima, 1937).
S. sibirica Lederer, 1853
 阿爾泰山 (Klima, 1937)
S. stötzneri Caradja, 1927
 四川 (Klima, 1937).
S. submedinella Caradja, 1927
 華西 (Klima, 1937).
S. tibetalis Caradja, 1937
 雲南 (Caradja, 1937).
S. truncicolella (Stainton, 1849)
 華西等 (Caradja, 1925; Klima, 1937).
S. vtrnotinctalis Hampson, 1896
 華中、華西 (Caradja, 1925; Klima, 1937).
S. xanthomelas De Joannis 1929
 廣東等 (Caradja & Meyrick, 1934; Klima, 1937).

擬螟亞科 *Pyraustinae* Ragonot 1891

- Adena mirifica* Caradja,
 廣東 (Caradja & Meyrick, 1934).
Aetholix falvibasalis (Guenée, 1859)
 華南 (Klima, 1937); 廣東等 (Caradja, 1925; Caradja & Meyrick, 1934).
Agathodes caliginodalis Snellen
 江西、西藏 (Caradja, 1925).
A. ostentalis (Hübner & Geyer, 1837)
 華東 (Caradja & Meyrick, 1934); 華西、台灣 (Shibuya, 1928); 福建、廣州 (Caradja, 1925).
Agrotera basinotata Hampson, 1891
 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).
Ag. discinotata Swinhoe, 1894
 華南 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
Ag. leucostola Hampson, 1896

華中 (Caradja, 1925; Klima, 1939); 華南 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 福建、廣州 (Caradja, 1925).

Ag. lienpingialis Caradja, 1925

華南 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Ag. nemoralis (Scopoli, 1763)

華北、華西 (Klima, 1939); 華中 (Caradja, 1925; Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939).

Ag. ornata (Wileman & South, 1917)

台灣 (Shibuya, 1928; Klima, 1939).

Ag. scissalis (Walker, 1865)

江西 (Caradja, 1925); 台灣 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Antigastra catalaunalis (Duponchel, 1833)

華中、上海 (Caradja, 1925); 台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

Archernis capitalis (Fabricius, 1794)

江西、福建、廣州 (Caradja, 1925); 台灣 (Caradja, 1925; Caradja & Meyrick, 1934; Shibuya, 1928 (*Protonoceras*)); 廣東 (Caradja & Meyrick, 1934).

Ar. humilis Swinhoe

浙江、福州、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Ar. tropicalis Walker

台灣 (Hampson, 1896; Caradja & Meyrick, 1934); 廣東 (Caradja & Meyrick, 1934).

Batyodes astalis Guenée, 1854

台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).

B. caldusalis (Walker, 1859)

華南 (Klima, 1939); 華西 (Caradja, 1925; Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣州 (Caradja, 1925).

B. interruptalis Caradja & Meyrick, 1934

廣東 (Caradja & Meyrick, 1934; Klima, 1939).

B. patulalis (Walker, 1865)

華南、華西 (Klima, 1939); 廣東 (Caradja, & Meyrick, 1934); 廣州 (Caradja, 1925).

B. principalis Leech, 1889

華中 (Caradja, 1925; Klima, 1939); 華南 (Klima, 1939); 華西 (Caradja, 1925; Klima, 1939); 江西、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).

Calamochrous acutellus Ev.

浙江 (Caradja, 1925).

C. dichroma Moore

廣州等 (Caradja, 1925).

C. eumorphalis Caradja, 1925

浙江 (Caradja, 1925; Caradja & Meyrick, 1934); 江西 (Caradja, 1925); 廣東、長江流域 (Caradja & Meyrick, 1934).

C. medialis Caradja, 1925

- 廣州 (Caradja, 1925).
- C. minimalis* Caradja, 1931
四川 (Caradja, 1931).
- C. obscuralis* Caradja, 1925
福建 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- C. pyraustalis* Strand, 1918
台灣 (Shibuya, 1928).
- C. tranquilalis* (Lederer, 1863)
台灣等 (Shibuya, 1928).
- Ceratarcha umbrosa* Swinhoe, 1894
華南 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).
- Chalcidoptera rufilinealis* Swinhoe, 1863
華南 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Cnaphalocroctis medinalis* (Guenée, 1854).
華南、華西 (Klima, 1939); 華中 (Caradja, 1925; Klima, 1939); 浙江、廣州、上海 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 江蘇、湖南、四川、廣西、江西 (李鳳藻, 1952).
- Cn. nursialis* (Walker, 1859)
廣東 (Caradja & Meyrick, 1934 (f.); Klima, 1939).
- Crocitolomia binotalis* Zeller, 1852
台灣 (Hampson, 1896; Shibuya, 1928; 李鳳藻, 1952).
- Cr. suffusalis* Hampson
江西 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).
- Cr. suffusalis* var. *sinensis* Caradja, 1925
廣州 (Caradja, 1925).
- Crocidophora artatalis* Caradja, 1925
福建、廣東 (Caradja, 1925).
- Cro. aurealis* Leech
浙江、福建 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- Cro. aurimargo* Warren
浙江 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- Cro. butleri* South
浙江 (Caradja, 1925).
- Cro. cuprotinctalis* Caradja, 1932
華南 (Caradja, 1932).
- Cro. discolorata* Swinhoe.
廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- Cro. evenoralis* (Walker, 1859)
華東、華中、華西、浙江、上海 (Caradja, 1925); 台灣 (Shibuya, 1928).
- Cro. flaviciliata* Hampson, 1910)

台灣 (Shibuya, 1928).

Cro. gensanalis South

廣東 (Caradja & Meyrick, 1934).

Cro. gladialis Leech

華中、華西、浙江、福建、廣州 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Cro. habisalis (Walker, 1859)

台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

Cro. heterogenalis Bremer

華中 (Caradja, 1925).

Cro. hoenei Caradja, 1932

華南 (Caradja, 1932).

Cro. limbolalis Moore

福州 (Caradja, 1925).

Cro. mectariphila Strand, 1918

台灣 (Shibuya, 1928).

Cro. nycterina Butler,

華中、浙江、廣東 (Caradja, 1925).

Cro. obscuralis South

華西 (Caradja, 1925).

Cro. pallida (Moore, 1888)

華中、福建、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Cro. rufitinctalis Hampson, 1918

台灣 (Shibuya, 1928).

Cro. tiermushana Caradja, 1935

華南 (Caradja, 1935).

Cro. zonalis Caradja, 1925

福建、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Diasemia accalis (Walker, 1859)

上海 (Hampson, 1896; Caradja, 1925); 華中、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

D. accalis var. *grammalis* Doubleday

華南、廣東、廣州 (Caradja, 1925).

D. calcaralis Strand, 1918

台灣 (Shibuya, 1925).

D. lepidoneuralis Strand, 1918

台灣 (Shibuya, 1928).

D. litterata Scopoli

華中、浙江、上海、台灣、廣東 (Caradja, 1925).

D. impulsalis var. *taiwanalis* Shibuya, 1928

台灣 (Shibuya, 1928).

D. ramburialis (Duponchel, 1832)

台灣等 (Shibuya 1928).

Diastictis acamasalis (Walker, 1859)

華中、華南 (Klima, 1939); 江西、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).

Di. adipalis (Lederer, 1863)

台灣等 (Shibuya, 1928; Klima, 1939).

Di. adipalis nigromarginalis Caradja, 1925

華南 (Klima, 1934); 廣州 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Di. aptalis (Walker, 1865)

台灣等 (Shibuya, 1928; Klima, 1939).

Di. aptalis kuangtungialis Caradja, 1925

華南 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Di. artificialis (Lederer, 1863)

台灣等 (Shibuya, 1928; Klima, 1939).

Di. fatualis (Lederer, 1863)

廣東 (Caradja & Meyrick, 1934).

Di. inspersalis (Zeller, 1852)

華北、華西 (Klima, 1939); 華東、華中 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Di. onychinalis (Guenée, 1854)

華南 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Di. telphusalis (Walker 1859)

台灣等 (Shibuya, 1928; Klima, 1939).

Dichocrocis chlorophanta (Butler, 1878)

台灣等 (Shibuya, 1928; Klima, 1939); 浙江 (李鳳藻, 1952).

Dicho. defnita (Butler, 1889)

華南 (Klima, 1939); 華中、華西 (Caradja, 1925; Klima, 1939); 廣東 (Caradja, & Meyrick, 1934); 廣州 (Caradja, 1925).

Dicho. diminutiva (Warren, 1896)

華南 (Klima, 1939); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Dicho. latipunctalis Caradja, 1925

華南 (Klima, 1939); 福建 (Caradja & Meyrick, 1934); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Dicho. nilusalis (Walker, 1859)

華南 (Klima, 1939); 華東、廣州 (Caradja, 1925); 華中、華西 (Caradja, 1925; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Dicho. plutusalis (Walker, 1859)

廣東等 (Caradja & Meyrick, 1934; Klima, 1939).

Dicho. punctiferalis (Guenée, 1854)

華中、華南、華西 (Klima, 1939); 浙江、上海 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934); 河北、陝西、江蘇、湖南、四川 (李鳳藻, 1952).

Dicho. rigidalis (Snellen, 1890)

廣東等 (Caradja & Meyrick, 1934; Klima, 1939).

Dicho. surusalis (Walker, 1859)

華中 (Caradja, 1925; Klima, 1939; 華南 (Klima, 1939); 台灣 (Shibuya, 1928; Caradja & Meyrick, 1934; Klima, 1939); 浙江、廣東 (Caradja & Meyrick, 1934).

Dicho. tigrina (Moore, 1886)

台灣等 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934 (f.)).

Dicho. zebalis (Moore, 1867)

華南 (Klima, 1939); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Discothyris ferruginata (Moore, 1888)

台灣等 (Shibuya, 1928).

Dis. megalophalis Hampson

廣州等 (Caradja, 1925).

Drosophantis corusca Meyrick, 1935

台灣 (Meyrick, 1935).

Eptpagis setinalis Hampson, 1918

台灣 (Shibuya, 1928).

Eptparbattia gloriosalis Caradja, 1925

福建 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Erecta chalybæis (Meyrick, 1885)

台灣等 (Shibuya, 1928 (*Hydriris*); Klima, 1939)

E. elutalis (Walker, 1859)

廣東等 (Caradja & Meyrick, 1934; Klima, 1939).

E. ornatalis (Duponchel, 1831).

華南 (Klima, 1939); 台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Euclasta defamatalis (Walker, 1859)

浙江、上海、福建、廣州 (Caradja, 1925), 台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

Eu. splendidalis Herr.-Schäff.

華中、華西等 (Caradja, 1925).

Eurrhyarodes bracteolalis (Zeller, 1852)

中國 (Hampson, 1896); 華中、華南、華西 (Klima, 1939); 上海、廣州 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Eur. controtalis (Hampson, 1900)

華南 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939).

Eur. leechi South, 1901

華北、華南 (Klima, 1939); 華中 (Caradja, 1925); 華西 (Caradja, 1925; Klima, 1939); 廣東

(Caradja, 1925; Caradja & Meyrick, 1934).

Eur. nymphulalis Strand, 1918

台灣 (Shibuya, 1928; Klima, 1939).

Eur. tricoloralis (Zeller, 1852)

華南 (Klima, 1939); 福建、廣州 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Evergestis extimalis Scopoli

華西 (Caradja, 1925); 華北 (李鳳蓀, 1952).

Ev. hoenei Caradja, 1939

長江流域 (Caradja, 1939).

Ev. hoenei f. ferruginea Caradja, 1939

長江流域 (Caradja, 1939).

Ev. juncalis Warren

華西 (Caradja, 1925).

Ev. orientalis gigantea Caradja, 1939

長江流域 (Caradja, 1939).

Exeristis minuscula Caradja, 1931

四川 (Caradja, 1931).

Ex. mystica Caradja, 1927

中國 (Caradja, 1927).

Filodes fulvidorsalis (Geyer, 1832)

廣東等 (Caradja & Meyrick, 1934; Klima, 1939).

F. mirificalis (Lederer, 1863)

華南 (Klima, 1939); 廣東 (Caradja & Meyrick, 1934 (var)).

Glyphodes actorionalis (Walker, 1859)

福建、南京、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928 (*Margaronia*)).

G. actorionalis var. *jovialis* (Felder & Rogenhofer, 1874)

廣東 (Caradja & Meyrick, 1934).

G. albifuscalis Hampson

華中、華西 (Caradja, 1925).

G. annulata (Fabricius, 1794)

華中、福建、廣州 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 台灣 (Shibuya, 1928 (*Margaronia*)).

G. badialis Walker

廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

G. bicolor var. *parvalis* Walker

廣東 (Caradja & Meyrick, 1934).

G. biotralis (Guenée, 1854)

台灣 (Hampson, 1896; Shibuya, 1928 (*Margaronia*)); 福建、南京、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

G. brizoalis Walker

華東 (李鳳藻, 1952).

G. caesalis Walker

福建 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

G. canthusalis (Walker, 1859)

台灣 (Hampson, 1896; Caradja, 1925; Shibuya, 1928 (*Margaronia*); Caradja & Meyrick, 1934);
廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

G. confinis Hampson, 1898

台灣等 (Hampson, 1898; Shibuya, 1928 (*Margaronia*)).

G. crithealis (Walker, 1859)

中國 (Hampson, 1896); 華中、華西、福建、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928 (*Margaronia*)); 廣東、雲南 (Caradja & Meyrick, 1934).

G. crithealis var. *minoralis* Caradja, 1925

華中、華西、福建 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

G. crithealis var. *parvipunctalis* Caradja, 1925

福建、台灣 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

G. eurytusalis (Walker, 1859)

台灣 (Shibuya, 1928 (*Margaronia*)).

G. fraterna Moore

福建 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

G. indica (Saunders, 1851)

華南、江蘇、四川 (李鳳藻, 1952); 浙江、福建、廣州 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928 (*Margaronia*)); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

G. itysalis (Walker, 1859)

台灣等 (Hampson, 1896; Shibuya, 1928 (*Margaronia*)).

G. marginata (Fabricius, 1794)

台灣 (Shibuya, 1928 (*Margaronia*)); 廣東 (Caradja & Meyrick, 1934).

G. melonella Linnaeus

河北、江蘇、浙江、廣東 (李鳳藻, 1952).

G. negatalis (Walker, 1859)

台灣 (Shibuya, 1928 (*Margaronia*)); 廣東 (Caradja & Meyrick, 1934).

G. nigribasalis Caradja, 1925

浙江 (Caradja, 1925).

G. nigropunctalis (Bremer, 1864)

台灣 (Shibuya, 1928 (*Margaronia*)); 東北六省 (李鳳藻, 1952).

G. perspectalis Walker

中國 (Hampson, 1896); 華東、華中、浙江、上海、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934); 江蘇、我國東部及沿海 (李鳳藻, 1952).

G. pfeifferae (Lederer, 1863)

台灣等 (Shibuya, 1928 (*Margaronia*)).

G. pomonalis Guenée

中國 (Hampson, 1896; Caradja, 1925).

G. pryeri Butler

廣東、四川 (Caradja & Meyrick, 1934); 浙江 (李鳳蓀, 1952).

G. pryeri var. *formosana* (Shibuya, 1928)

台灣 (Shibuya, 1928 (*Margaronia*)).

G. pulverulentalis Hampson, 1896

浙江、海南 (Caradja, 1925).

G. pyloalis (Walker, 1859)

中國 (Hampson, 1896); 華東、華中、華西、浙江、福建、上海 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928 (*Margaronia*)); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 江蘇、四川、沿海諸省 (李鳳蓀, 1952).

G. quadrimaculalis Bremer

華西、浙江 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

G. quinquepunctalis Hubner

廣東 (Caradja & Meyrick, 1934).

G. stolalis (Guenée, 1854)

華西、台灣 (Shibuya, 1928 (*Margaronia*)).

G. tienmushana Caradja, 1935

華南 (Caradja, 1935)

G. vertumnalis Guenée

廣東 (Caradja & Meyrick, 1934).

G. warrenalis Swinhoe, 1894

台灣 (Shibuya, 1928 (*Margaronia*)).

Goniorhynchus butyrosa (Butler, 1879)

華中 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).

Go. marginalis Warren, 1895

中國 (Klima, 1939).

Hapalia teneralis Caradja, 1939

長江流域 (Caradja, 1939).

Heterocnephes lubricosa (Warren, 1896)

華中、華南 (Klima, 1939); 上海、廣州 (Caradja, 1925); 浙江、廣東 (Caradja & Meyrick, 1934).

Heliothela cretostrigalis Caradja, 1925

廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

H. nigralbata Hampson, 1899

中國 (Hampson, 1899); 華東 (Caradja, 1925).

Hellula angarensis Caradja, 1939

山西 (Caradja, 1939).

He. undalis (Fabricius, 1794)

台灣 (Shibuya, 1928 (*Oebia*)); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925); 湖南、湖北、四川 (李鳳蓀, 1952).

Hemiscopsis cinerea Warren, 1892

台灣等 (Shibuya, 1928).

Hem. expansa Warren

廣東 (Caradja & Meyrick, 1934).

Hem. suffusalis Walker

上海、台灣 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Heortia vitessoides Moore

香港 (Hampson, 1896; Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Hyalobathra coenostolalis (Snellen, 1890)

台灣 (Shibuya, 1928); 廣東、廣州 (Caradja, 1925).

Hy. coenostolalis var. *udeoides* (Hampson, 1891)

廣東 (Caradja & Meyrick, 1934).

Hy. filalis (Guenée, 1854)

台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

Hy. illectalis (Walker, 1859)

浙江 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

Hy. intermedialis Caradja 1939

陝西 (Caradja, 1939).

Hy. micralis Caradja, 1932

華南 (Caradja, 1932).

Hy. opheltialis Walker

廣東等 (Caradja, & Meyrick, 1934).

Hy. undulinea (Hampson, 1891)

台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934 (*aequalis*)); 廣州 (Caradja, 1925 (*undulinea*)).

Hymenja perspectalis (Hübner, 1796)

台灣等 (Shibuya, 1928; Klima, 1939).

Hym. recurvalis (Fabricius, 1775)

華中、廣東 (Caradja, 1925); 華南、華西 (Klima, 1939); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1939); 江西、我國沿海諸省 (李鳳藻, 1952).

Hyperanalyta pseudoapicalis (Strand, 1918)

台灣 (Caradja, 1925; Shibuya, 1928 (*Leucinodes apicalis*)); Caradja & Meyrick, 1934 (*L. apicalis*); Strand, 1918); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Ischnurges gratosalis (Walker, 1859)

華北 (Hampson, 1896); 華南、福建、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

I. minoralis Caradja, 1931

廣州 (Caradja, 1931).

I. octoguttalis Felder & Eögenhofer, 1874

台灣等 (Shibuya, 1928).

Lamprohata mirabilis Caradja, 1925

廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Lamprosema annubilata (Swinhoe, 1894)

廣東 (Caradja, 1925; Klima, 1939).

L. bianoralis (Walker, 1859)

台灣等 (Shibuya, 1928; Klima, 1939).

L. biformis (Butler, 1889)

台灣 (Shibuya, 1928; Klima, 1939).

L. catenalis (Wileman, 1911)

華西、華南 (Klima, 1939); 台灣 (Caradja & Meyrick, 1934); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

L. charæsalis (Warren, 1859)

華南 (Klima, 1939); 台灣 (Caradja, 1925; Shibuya, 1928; Caradja & Meyrick, 1934; Klima, 1939); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

L. chrysorycta (Meyrick, 1884)

華西 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939).

L. commixta (Butler, 1879)

華中 (Caradja, 1925); 華南、華西 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

L. cyanealis (Walker, 1859)

華南 (Klima, 1939); 江西 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

L. dairalis (Walker, 1859)

華北、華南、華西 (Klima, 1939); 廣東 (Caradja & Meyrick, 1934).

L. diemenalis (Guenée, 1854)

華南 (Klima, 1939); 華西 (Caradja, 1925; Klima, 1939); 台灣 (Hampson, 1896 (*Nacoleia*); Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

L. eximinalis (Warren, 1896)

華南 (Klima, 1939).

L. fusalis (Warren, 1896)

華中 (Caradja 1925; Klima, 1939).

L. hoenei Caradja, 1932

華南 (Caradja, 1932; Klima, 1939).

L. immundalis (South, 1901)

華中 (Caradja, 1925); 湖北 (Klima, 1939).

L. indicata (Fabricius, 1775)

華北、華南、華西 (Klima, 1939); 華中 (Caradja, 1925; Klima, 1939); 浙江、上海、福建、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja 1925; Caradja & Meyrick, 1934).

L. invalidalis (South, 1901)

華中、華西等 (Klima, 1939).

L. karenkonalis Shibuya, 1928

台灣 (Shibuya, 1938; Klima, 1939).

L. marionalis (Walker, 1859)

華南、華西 (Klima, 1939); 華中 (Caradja, 1925; Klima, 1939); 廣東 (Caradja, 1925; Caradja &

Meyrick, 1934).

L. minoralis (Walker, 1885)

華南、華西 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

L. misera (Butler, 1879)

中國 (Klima, 1939).

L. ochrimaculalis (South, 1910)

華南 (Klima, 1939); 華中 (Caradja, 1925; Klima, 1939); 福建 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

L. pallidnotalis (Hampson, 1912)

華中、華西 (Caradja, 1925; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).

L. pectinalis Hampson, 1918

華中 (Klima, 1939).

L. perstygialis Hampson, 1912

華中 (Klima, 1939); 台灣、廣東 (Caradja & Meyrick, 1934).

L. perstygialis Hampson, 1912

華南 (Klima, 1939); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

L. poconalis (Walker, 1859)

華中、華南、華西 (Klima, 1939).

L. poconalis f. *decisalis* (Walker, 1865)

華中 (Caradja, 1925; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

L. sauteplialis (Strand, 1918)

台灣 (Shibuya, 1928; Klima, 1939).

L. sibirialis (Millière, 1879)

華中、華西 (Caradja, 1925; Klima, 1939); 浙江 (Caradja, 1925); 廣東、四川 (Caradja & Meyrick, 1934).

L. similis (Moore, 1885)

中國 (Klima, 1939); 台灣、廣東 (Caradja & Meyrick, 1934).

L. subbalis (Hampson, 1912)

華西 (Klima, 1939).

L. subargentalis (Snellen, 1890)

華西等 (Caradja, 1925; Klima, 1939).

L. stigmatialis (Warren, 1896)

華南 (Klima, 1939); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

L. tampusalis (Walker, 1859)

華南 (Klima, 1939); 江西、福建 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

L. tienmushanus Caradja, 1935)

華南 (Caradja, 1925; Klima, 1939).

L. tristrialis (Bremer, 1864)

華中 (Caradja, 1925; Klima, 1939); 華西 (Klima, 1939); 上海 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).

Lepidoneura longipalpis (Swinhoe, 1894)

台灣 (Shibuya, 1928).

Lepyrodes geometralis Guenée, 1854

台灣 (Hampson, 1896; Caradja, 1925; Shibuya, 1928 (*Nausinoe*)); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Le. perspectata Fabricius

福建、廣州 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928 (*Nausinoe*)); 廣東 (Caradja & Meyrick, 1934).

Leucinodes orbonalis Guenée, 1854

華南 (李鳳蓀, 1952); 台灣 (Shibuya, 1928; 李鳳蓀 1952).

Leu. perlucidalis Caradja & Meyrick, 1934

廣東 (Caradja & Meyrick, 1934).

Loxostege anpingialis (Strand, 1918)

台灣 (Shibuya, 1928).

Lox. concoloralis Lederer

廣州等 (Caradja, 1925 (*Phlyctaenodes*))

Lox. confusalis South

華西 (Caradja, 1925 (*Phlyctaenodes*)).

Lox. decoloralis sinensis (Caradja, 1925)

上海 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934 (*Phlyctaenodes*)).

Lox. diaphana (Caradja & Meyrick, 1934)

浙江、廣東 (Caradja & Meyrick, 1934 (*Phlyctaenodes*)).

Lox. formosibia (Strand, 1918)

台灣 (Shibuya, 1928).

Lox. massalis (Walker, 1859)

台灣等 (Shibuya, 1928).

Lox. palealis Schiff.

中國 (Hampson, 1896 (*Phlyctaenodes*)); 華中、上海 (Caradja, 1925 (var.)); 廣東 (Caradja & Meyrick, 1934 (*Phlyctaenodes*)).

Lox. sticticalis tenebrosa Caradja, 1939

長江流域 (Caradja, 1939).

Lox. sulphuralis f. *minor* Caradja, 1937

中國 (Caradja, 1937).

Lox. tschelialis Caradja, 1927

北京 (Caradja, 1927).

Lox. umbrosalis Warren

華中、華西 (Caradja, 1925 (*Phlyctaenodes*)); 廣東 (Caradja & Meyrick, 1934 (*Phlyctaenodes*)).

Lox. verticalis Linnaeus

華中、華西 (Caradja, 1925 (*Phlyctaenodes*)).

Lox. yuennanensis Caradja, 1937

雲南 (Caradja, 1937).

Lygrotia amynusalis Walker

廣東等 (Caradja & Meyrick, 1934).

Ly. euryclealis Walker

浙江、台灣等 (Caradja, 1925).

Ly. euryclealis minutalis Caradja & Meyrick, 1934

廣東等 (Caradja & Meyrick, 1934).

Ly. obrinusalis Walker

廣東等 (Caradja & Meyrick, 1934).

Ly. obrinusalis teneralis Caradja, 1932.

華南 (Caradja, 1932).

Ly. quaternalis (Zeller, 1852)

華中、華西、廣州 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928).

Ly. quaternalis f. *delicatilis* (Hampson)

華中、華西、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Ly. syleptalis (Strand, 1918)

台灣 (Shibuya, 1928).

Marasmia bilinealis Hampson, 1891

華西 (Klima, 1939).

M. bilinealis intristalis Caradja, 1927

華西 (Klima, 1939).

M. euryterminalis Hampson, 1917

台灣等 (Shibuya, 1928; Klima, 1939).

M. latimarginalis Hampson, 1891

華南 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

M. pauperalis Strand, 1918

台灣 (Shibuya, 1928; Klima, 1939).

M. trapezalis (Guenée, 1854)

華南 (Klima, 1939); 福建、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

M. trebiusalis (Walker, 1859)

華南 (Klima, 1939).

M. venilialis (Walker, 1859)

華南、華西 (Klima, 1939); 台灣 (Caradja, 1925; Shibuya, 1928); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Maruca amboinensis (Felder & Rogenhofer, 1874)

台灣 (Caradja, 1925; Shibuya, 1928); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925 (*Crochiphora*)).

Ma. testulalis (Geyer, 1832)

華南 (李鳳蓀, 1952); 華中、華西、浙江、上海、廣州 (Caradja, 1925 (*Crochiphora*)); 台灣 (Shibuya, 1928; 李鳳蓀, 1952); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Massepha absolutalis Walker, 1859

台灣 (Shibuya, 1928).

Mecyna polygonalis var. *carbonalis* Caradja, 1939

雲南 (Caradja, 1939).

Meroctena tullalis Walker

廣東 (Caradja & Meyrick, 1934).

Metasia coniotalis Hampson, 1903

台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

Me. hodiusalis Walker

中國 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Me. morbidalis South, 1901

中國 (South, 1901); 華西 (Caradja, 1925).

Me. paganalis South, 1901

中國 (South, 1901); 華東、浙江 (Caradja, 1925).

Me. vicinalis South, 1901

中國 (South, 1901); 華中、華西 (Caradja, 1925).

Nacoleia cinisalis Caradja, 1937

中國 (Caradja, 1937).

N. immundalis South, 1901

中國 (South, 1901).

N. maculalis South, 1901

中國 (South, 1901).

N. ochrimaculalis South, 1901

中國 (South, 1901).

Neurina procopia (Stoll, 1781)

台灣 (Shibuya 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).

Nomophila noctuella (Schifferrmüller & Denis, 1775)

中國 (Caradja, 1925); 台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

Noorda amethystina (Swinhoe, 1894)

台灣等 (Shibuya, 1928).

No. ignealis Hampson, 1899

台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

Nosophora albigitallis Swinhoe, 1890

華南 (Klima, 1939); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Nos. chironalis (Walker, 1859)

華中 (Klima, 1939); 上海 (Caradja, 1925).

Nos. dispilalis Hampson, 1898

廣東等 (Caradja & Meyrick, 1934).

Nos. dispilalis artusalis Caradja, 1925

華南 (Klima, 1939); 廣州 (Caradja, 1925).

Nos. semitritalis (Lederer, 1863)

華中、華南 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934);

福建 (李鳳藻, 1952).

Nos. semitratalis hilaralis Caradja, 1925

華中 (Caradja, 1925; Klima, 1939); 福建、廣東、廣州 (Caradja, 1925).

Nos. semitratalis var. *orbicularis* Shibuya, 1928

台灣等 (Shibuya, 1928; Klima, 1939).

Nos. taihoqualis Strand, 1918

台灣 (Shibuya, 1928; Klima, 1939).

Omphisa albalis Caradja, 1925

青島、浙江、上海 (Caradja, 1925).

O. anastomosalis Guenée

中國 (Hampson, 1896); 華南、台灣、海南 (李鳳藻, 1952).

O. illisalis (Walker, 1859)

華南 (李鳳藻, 1952); 福建、廣州 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

O. plagialis

陝西 (李鳳藻, 1952).

Orthoraphis bilineata Caradja

廣東 (Caradja & Meyrick, 1934).

Pagya amphisalis (Walker, 1859)

華中、華西 (Caradja, 1925; Klima, 1939); 華南 (Klima, 1939); 福建 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

P. arbiter (Butler, 1879)

台灣等 (Shibuya, 1928; Klima, 1939).

P. auroralis (Moore, 1887)

華南 (Klima, 1939); 江西、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

P. botydalis (Snellen, 1880)

台灣等 (Shibuya, 1928; Klima, 1939).

P. discolor Swinhoe, 1894

廣東等 (Caradja & Meyrick, 1934; Klima, 1939).

P. griseotincta Caradja, 1939

長江流域 (Caradja, 1939).

P. lustralis Snellen, 1890

華南 (Klima, 1939); 江西 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

P. nebulosa Wileman & South, 1917

台灣 (Shibuya, 1928; Klima, 1939).

P. pullalis Swinhoe, 1903

廣東等 (Caradja & Meyrick, 1934; Klima, 1939).

P. quinquelineata Hering, 1903

華中、華西 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939).

P. salvalis (Walker, 1859)

華中 (Caradja, 1925; Klima, 1939); 華南、華西 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Paratalanta labutonalis amuralis Rom.

華中、華西、上海 (Caradja, 1925 (var.)); 廣東 (Caradja & Meyrick, 1934).

Pa. ussurialis (Bremer, 1864)

台灣等 (Shibuya, 1928).

Parbattia latifascialis South, 1901

中國 (South, 1901); 華中 (Caradja, 1925).

Par. vialis Moore, 1888

台灣等 (Shibuya, 1928).

Phostria analis (Snellen, 1880)

華南 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Ph. crithonalis (Walker, 1859)

華南 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Ph. discipunctalis (Hampson, 1903)

廣東等 (Caradja & Meyrick, 1934; Klima, 1939).

Ph. expictalis (Christoph, 1881)

華北 (Klima, 1939).

Ph. glyphodalis (Walker, 1865)

廣東等 (Caradja & Meyrick, 1934; Klima, 1939).

Ph. imbecilis Moore, 1887

華南 (Klima, 1939); 廣東 (Caradja & Meyrick, 1934).

Ph. maculicostalis (Hampson, 1893)

廣東等 (Caradja & Meyrick, 1934; Klima, 1939).

Ph. noctescens (Moore, 1888)

華南、華西 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Ph. obscurata (Moore, 1886)

台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

Ph. palliventralis (Snellen, 1880)

華北、華南、華西 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Ph. palliventralis var. *dilutior* Caradja

廣東 (Caradja & Meyrick, 1934)

Ph. selenalis Caradja & Meyrick, 1934

廣東 (Caradja & Meyrick, 1934; Klima, 1939).

Ph. unitalis (Guenée, 1854)

華南 (Klima, 1939); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Ph. unilinctalis (Hampson, 1896)

廣東等 (Caradja & Meyrick, 1934; Klima, 1939).

Pilocrocis contortalis Hampson

華南、華中、廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Pionea albifimbrialis (Walker, 1865)

華東 (Caradja, 1925,); 台灣 (Caradja, 1925; Shibuya, 1928 (*Udea*))

Pt. albopedalis Motschulsky

華中 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Pt. auratalis (Warren, 1895).

廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 台灣 (Shibuya, 1928 (*Udea*)).

Pt. auratalis ab. *obscura* Caradja, 1935

華南 (Caradja, 1935).

Pt. aureolalis f. *contractalis* Warren

廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Pt. brevialis (Walker, 1859)

江西 (Caradja, 1925); 台灣 (Shibuya, 1928 (*Udea*)); 廣東 (Caradja & Meyrick, 1934).

Pt. endotrichialis (Hampson, 1918)

台灣 (Shibuya, 1928 (*Udea*)).

Pt. fentoni Butler

華中 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Pt. ferrugalis (Hübner, 1796)

華東、華西、上海、廣東、廣州、(Caradja, 1925); 台灣 (Shibuya, 1928 (*Udea*)).

Pt. flavicinctalis Snellen

廣東 (Caradja, & Meyrick, 1934); 廣州 (Caradja, 1925).

Pt. flavicinctalis Snellen

廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Pt. flavofimbriata (Moore, 1888)

台灣 (Shibuya, 1928 (*Udea*)); 廣東 (Caradja & Meyrick, 1934).

Pt. forficalis Linnaeus

華北 (李鳳藻, 1952); 廣東 (Caradja & Meyrick, 1934).

Pt. genialis Lpech

華東 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Pt. incertalis Caradja, 1937

雲南 (Caradja, 1937).

Pt. ingentalis Caradja, 1925

廣東、廣州 (Caradja, 1925).

Pt. inornata Butler

廣東 (Caradja 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Pt. lolotialis Caradja, 1927

中國 (Caradja, 1927).

Pt. lolotialis f. *intunecalis* Caradja 1927

中國 (Caradja, 1927).

Pt. leucanalis Swinhoe

- 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 江西 (Caradja, 1925).
- Pi. lienpinglalis* Caradja, 1925
廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Pi. lugubralis* Leech
華中 (Caradja, 1925).
- Pi. mendicalis* South, 1901
中國 (South, 1901); 華中 (Caradja, 1925).
- Pi. minnehaha* (Pryer, 1877)
華中、上海 (Caradja, 1925 (*minnenhaha*)); 台灣 (Shibuya, 1928 (*Udea*)).
- Pi. nea* Strand, 1918
台灣 (Shibuya, 1928 (*Udea*)).
- Pi. nephelitalis* Hampson
浙江、湖北 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Pi. nigrostigmatis* Warren
廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).
- Pi. ochrealis* Moore
廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).
- Pi. orbicentralis* Chr.
華西等 (Caradja, 1925).
- Pi. pallidalis* South, 1901
中國 (South, 1901); 華中 (Caradja, 1925).
- Pi. pandalis* Hübner
華東、華中、華西、上海 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Pi. planalis* South, 1901
中國 (South, 1901); 華西 (Caradja, 1925).
- Pi. polistotalis* (Hampson, 1918)
台灣 (Shibuya, 1928 (*Udea*)).
- Pi. pseudocrocealis* South
廣東等 (Caradja & Meyrick, 1934).
- Pi. pseudotornata* Caradja, 1931
四川 (Caradja, 1931).
- Pi. punctiferalis* South, 1901
中國 (South, 1901); 華中 (Caradja, 1925).
- Pi. puralis* South, 1901
中國 (South, 1901); 華中 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934 (*pseudotornata*)).
- Pi. rubignalis* Hübner
廣東 (Caradja & Meyrick, 1934).
- Pi. sanguinalis priscalis* Caradja, 1935
華南 (Caradja, 1935)
- Pi. schaeferi* Caradja, 1937
雲南 (Caradja, 1937).

- Pi. stigmatalis* Wileman, 1911
台灣等 (Shibuya, 1928 (*Udea*)).
- Pi. subplanalis* Caradja, 1937
雲南 (Caradja, 1937).
- Pi. testacea* (Butler)
華東、華西、上海、廣州 (Caradja, 1925 (f.)); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Pi. thyalis* Walker
中國 (Caradja, 1925).
- Pi. tritalis* Christoph.
華南 (Caradja, 1925).
- Pi. verbascalis* Schiff ermillier.
廣東 (Caradja & Meyrick, 1934).
- Pi. verbascalis* f. *egentalis* Christoph.
廣東 (Caradja, 1925; Caradja & Meyrick, 1934 (sp.)).
- Pi. verbascalis* f. *plumbocillialis* Snellen
華東、華中、華西、上海 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934 (sp.)).
- Polygrammodes priscalis* Caradja, 1937
雲南 (Caradja, 1927).
- Po. sabelialis* (Guenée, 1854)
福建 (Caradja, 1925; Caradja & Meyrick, 1934 (*nigropygalis*)); 台灣 (Caradja, 1925; Shibuya, 1928; Caradja & Meyrick, 1934 (*nigropygalis*)); 廣東 (Caradja & Meyrick, 1934 (*nigropygalis*)); 廣州 (Caradja, 1925).
- Po. thoosalis* (Walker, 1859)
台灣 (Shibuya, 1928).
- Polythlipta divaricata* Moore, 1886
台灣等 (Shibuya, 1928).
- Poly. euroalis* Swinhoe
廣東 (Caradja & Meyrick, 1934).
- Poly. liquidalis* Leech
華中、華西 (Caradja, 1925); 廣東、四川 (Caradja & Meyrick, 1934).
- Poly. maculalis* South, 1901
中國 (South, 1901); 華中、福建、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- Procdema inscisale* (Walker)
廣東等 (Caradja & Meyrick, 1934).
- Prorodes mimica* Swinhoe, 1894
台灣等 (Shibuya, 1928; Klima, 1939).
- Psara aegrotalis* Zeller, 1852
浙江、江西、上海、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- Psara (Pachyzancla) aegrotalis* f. *rudia* (Warren)
中國 (Hampson, 1896).
- Ps. basalis* (Walker, 1865)

- 台灣等 (Shibuya, 1928).
- Ps. cynaralis* (Walker, 1859)
台灣等 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).
- Ps. dilatipes* (Walker, 1865)
台灣等 (Shibuya, 1928).
- Ps. elongalis* (Warren, 1880)
台灣 (Shibuya, 1928).
- Ps. hipponalis* (Walker, 1859)
台灣等 (Shibuya, 1928).
- Ps. intensalis* Swinhoe
江西、福建 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Ps. intensalis rubellalis* Caradja, 1925
江西、福建、廣東 (Caradja, 1925).
- Ps. licarsisalis* (Walker, 1859)
中國 (Hampson, 1896); 華中、江西、上海、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).
- Ps. marginalis* Warren
華中、華西、浙江、江西 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Ps. mimeticalis* (Hering, 1901)
台灣等 (Shibuya, 1928).
- Ps. phaeopteralis* Guenée
廣東 (Caradja & Meyrick, 1934).
- Ps. stultalis* Walker
中國 (Hampson, 1896 (*Pachyzancla*); Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- Ps. subalbescens* (Swinhoe, 1894)
台灣等 (Shibuya, 1928).
- Ps. submarginalis* Caradja, 1925
浙江 (Caradja, 1925).
- Ps. taiwanalis* Shibuya, 1928
台灣 (Shibuya, 1928).
- Pycnarmon aeriferalis* (Moore, 1877)
華南 (Klima, 1939); 江西 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Py. geminipuncta* Hampson, 1912
華西 (Klima, 1939).
- Py. cribrata* (Fabricius, 1794)
華南、華西 (Klima, 1939); 華中 (Caradja, 1925; Klima, 1939); 台灣 (Hampson, 1896 (*caberalis*); Caradja, 1925 (*caberalis*); Shibuya, 1928; Klima 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).
- Py. (Aripa), Pyralosymatophora frenulalis* Strand, 1918
台灣 (Strand, 1918).

- Py. jaguaralis* (Guenée, 1854)
華西等 (Klima, 1939).
- Py. jaguaralis* var. *chitensis* Leech, 1901
華西等 (Caradja, 1925; Klima, 1939).
- Py. lactiferalis* (Walker, 1859)
中國 (Hampson, 1896); 華南、華西 (Klima, 1939), 華中、廣州 (Caradja, 1925); 台灣 (Caradja, 1925; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Py. marginalis* (Snellen, 1890)
台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).
- Py. meritalis* (Walker, 1859)
中國 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934 (*plexippusalis*)).
- Py. pantherata* (Butler, 1878)
華中、華西 (Caradja, 1925; Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 浙江 (Caradja, 1925).
- Py. radiata* (Warren, 1896)
華西 (Caradja, 1925; Klima, 1939); 華中、華南 (Klima, 1939); 福建、廣東 (Caradja, 1925).
- Py. radiata benesignata* Caradja, 1925
廣東 (Caradja & Meyrick, 1934 (f.)); 華南 (Klima, 1939).
- Py. tylostegalis* (Hampson, 1900)
華中、華西 (Caradja, 1925; Klima, 1939); 華南 (Klima, 1939); 浙江、福建、上海 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- Pygospila incertalis* Caradja, 1937
雲南 (Caradja, 1937).
- Pygo. minoralis* Caradja, 1937
雲南 (Caradja, 1937).
- Pygo. tyres* Cramer
廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1924).
- Pygo. guennanensis* Caradja, 1937
雲南 (Caradja, 1937).
- Pyrausta aerealis* f. *glaucalis* Caradja, 1927
中國 (Caradja, 1927).
- Pyr. auranticillialis* Caradja, 1925
廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).
- Pyr. ausontalis* (Snellen)
廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 台灣 (Shibuya, 1928 (*deductalis*)).
- Pyr. austa* Strand, 1918
台灣 (Shibuya, 1928).
- Pyr. baibarensis* Shibuya, 1928
台灣 (Shibuya, 1928).
- Pyr. bielti* Oberthür

- 華西 (Caradja, 1925).
- Pyr. carminalis* Caradja, 1925
浙江 (Caradja & Meyrick, 1934); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Pyr. celatalis* (Walker, 1859)
台灣 (Hampson, 1896; Shibuya, 1928); 浙江 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- Pyr. cespitales* Schiff er Müller.
華中、華西、上海 (Caradja, 1925).
- Pyr. cespitalis yangtsealis* Caradja, 1938
雲南 (Caradja, 1938).
- Pyr. clausalis subcalusalis* Caradja & Meyrick, 1934
廣東 (Caradja & Meyrick, 1934).
- Pyr. coclesalis* (Walker, 1859)
華中、浙江、上海、廣州 (Caradja, 1925); 台灣 (Hampson, 1896; Caradja, 1925; Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).
- Pyr. contigualis* South
華西 (Caradja, 1925).
- Pyr. contristalis* Caradja, 1932
華南 (Caradja, 1932).
- Pyr. culminivola* Caradja, 1939
長江流域 (Caradja, 1939).
- Pyr. curealis* Leech
華東、華南 (Caradja, 1925).
- Pyr. damoalis* Walker
中國 (Hampson, 1896); 華中、華西、江西、上海 (Caradja, 1925).
- Pyr. delicatalis* South
華中、華西 (Caradja, 1925).
- Pyr. dintasalis* (Walker, 1859)
華中、華西、廣州 (Caradja, 1925); 上海 (Hampson, 1896; Caradja, 1925); 台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).
- Pyr. diniasalis* var. *capnosalis* Caradja, 1925
廣州 (Caradja, 1925).
- Pyr. discrimaculalis* Hampson
北京等 (Caradja, 1925).
- Pyr. distans* Moore
華中、華西 (Caradja, 1925).
- Pyr. draeskei* Caradja, 1927
中國 (Caradja, 1927).
- Pyr. erioptisalis* Walker
浙江等 (Caradja, 1925).
- Pyr. ferrifuscalis* Hampson
廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Pyr. flavalis (Schifferrmüller)

上海 (Caradja, 1925),

Pyr. fuscalis Schifferrmüller

上海等 (Caradja, 1925).

Pyr. fuscobrunnealis South

華中、華西 (Caradja, 1925).

Pyr. gentilis South

華西 (Caradja, 1925).

Pyr. gentilis reductalis Caradja, 1937

雲南 (Caradja, 1937).

Pyr. gracilis (Butler, 1879)

華南、華西 (Caradja, 1925 (*gracilalis*)); 台灣 (Shibuya, 1928).

Pyr. gracilis meridionalis Caradja & Meyrick, 1934

西南、雲南 (Caradja & Meyrick, 1934).

Pyr. griseocollalis South

華西 (Caradja, 1925).

Pyr. hampsoni South

華中、華西 (Caradja, 1925).

Pyr. hoozana Strand, 1918

台灣 (Shibuya, 1928).

Pyr. hyalodiscalis (Warren, 1895)

台灣等 (Shibuya, 1928).

Pyr. incoloralis (Guenée, 1854)

台灣等 (Shibuya, 1928).

Pyr. leechi South

華西 (Caradja, 1925).

Pyr. limbata Butler

華中、華西、上海 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).

Pyr. machoeratis (Walker, 1859)

台灣 (Hampson, 1896; Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).

Pyr. machoeratis rubicundalis (Warren, 1896)

台灣 (Caradja, 1925); Shibuya, 1928 (sp.); 廣東 (Caradja & Meyrick, 1934 (sp.)); 廣州 (Caradja, 1925).

Pyr. maenialis Oberthür

華西 (Caradja, 1925).

Pyr. mandarinialis South

華西 (Caradja, 1925).

Pyr. masculina Strand, 1918

台灣 (Shibuya, 1928).

Pyr. minimalis Caradja, 1925

廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Pyr. moderatilis Christoph

華中、華西 (Caradja, 1925); 廣東、雲南 (Caradja & Meyrick, 1934).

Pyr. moupinialis South

上海 (Caradja, 1925).

Pyr. mundalis South

華中 (Caradja, 1925).

Pyr. mystica Caradja, 1932

華南 (Caradja, 1932).

Pyr. nubilalis Hübner

華中、華西、上海 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934); 河北、陝西、察哈爾、東北六省、山東、江蘇、浙江、台灣、河南、湖南、四川、(李鳳蓀 1952).

Pyr. oberthürti South

華西 (Caradja, 1925).

Pyr. obliquata (Moore, 1888)

台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934 (*obliqua*)).

Pyr. obscurior Caradja, 1938

雲南 (Caradja, 1938).

Pyr. ochracealis Walker

廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Pyr. octomaculalis Linnaeus

上海等 (Caradja, 1925).

Pyr. odontogrammalis Caradja, 1925

廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).

Pyr. palustralis infascialis Caradja, 1925

江西、上海、廣東 (Caradja, 1925).

Pyr. pala Strand, 1918

台灣 (Shibuya, 1928).

Pyr. perstimilis Caradja, 1932

華南 (Caradja, 1932).

Pyr. phoenicealis (Hübner, 1818)

中國 (Hampson, 1896); 浙江、福建 (Caradja, 1925); 台灣 (Shibuya, 1928).

Pyr. postalbalis South, 1901

中國 (South, 1901); 華西 (Caradja, 1925).

Pyr. punctilinealis South

華西 (Caradja, 1925).

Pyr. pygmaealis South

華中 (Caradja, 1925).

Pyr. quadrimaculalis South

華西 (Caradja, 1925).

Pyr. rubellalis (Snellen, 1890)

台灣 (Shibuya, 1928); 浙江、廣東 (Caradja & Meyrick, 1934).

- Pyr. rufalis* South
華西 (Caradja, 1925).
- Pyr. salentialis* Snellen
廣東 (Caradja & Meyrick, 1934).
- Pyr. sericealis* (Wileman & South, 1917)
台灣 (Shibuya, 1928).
- Pyr. signalalis* Walker
華中 (Caradja, 1925).
- Pyr. sikktma* Moore
華中、華西 (Caradja, 1925).
- Pyr. sclaris* Caradja, 1938
雲南 (Caradja, 1938).
- Pyr. splendida* Caradja, 1939
雲南 (Caradja, 1939).
- Pyr. stötzneri* Caradja, 1927
中國 (Caradja, 1927).
- Pyr. strigatalis* Caradja, 1937
雲南 (Caradja, 1937).
- Pyr. suavis* Walker
廣東 (Caradja & Meyrick, 1934).
- Pyr. subcrocealis* (Snellen)
廣東 (Caradja & Meyrick, 1934).
- Pyr. subfuscalis* Caradja & Meyrick, 1934
廣東 (Caradja & Meyrick, 1934).
- Pyr. sumptuosalis* Caradja, 1927
中國 (Caradja, 1927).
- Pyr. syfianialis* Oberthür
華西 (Caradja, 1925).
- Pyr. szetschuanalis* Caradja, 1927
中國 (Caradja, 1927).
- Pyr. tamsi* Caradja, 1927
中國 (Caradja, 1927).
- Pyr. tapashanensis* Caradja, 1939
陝西 (Caradja, 1939).
- Pyr. tibetalis* Oberthür
華西 (Caradja, 1925).
- Pyr. tithonalis* Zeller
華西等 (Caradja, 1925).
- Pyr. tithonalis* var. *latiplagiatis* Caradja, 1935
中國 (Caradja, 1935).
- Pyr. topa* Strand, 1918

- 台灣 (Shibuya, 1928).
- Pyr. tortualis* South
- 華西 (Caradja, 1925).
- Pyr. triplagalis* Warren
- 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 四川 (Caradja & Meyrick, 1934).
- Pyr. unipunctata* Butler
- 華中、華西 (Caradja, 1925).
- Pyr. vicinalis* South, 1901
- 華中 (Caradja, 1925); 台灣 (Shibuya, 1928).
- Pyr. vinacealis* Caradja, 1925
- 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Ravanod xiphialis* Walker, 1859
- 廣東 (Caradja & Meyrick, 1934).
- Rehimena phrynealis* (Walker, 1859)
- 華中 (Caradja, 1925; Klima, 1939); 華西 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).
- R. reductalis* Caradja, 1932
- 上海 (Caradja, 1932; Klima, 1939).
- R. straminealis* South, 1901
- 中國 (South, 1901); 華中 (Caradja, 1925; Klima, 1939).
- Sameodes bistigmalis* Fryer
- 華東、華中、浙江 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- S. cancellalis* (Zeller, 1852)
- 福建 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928 (*Epipagis*)); 廣東 (Caradja & Meyrick, 1934).
- S. miltochristalis* Hampson, 1896
- 台灣 (Shibuya, 1928 (*Epipagis*)); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Simaethis (Simaethitis) leechi* South, 1901
- 中國 (South, 1901); 華西 (Caradja, 1925).
- Sylepta adductalis* Walker
- 廣東等 (Caradja & Meyrick, 1934).
- Sy. amoyalis* Caradja, 1925
- 福建 (Caradja, 1925; Klima, 1939).
- Sy. angustimaculalis* Shibuya, 1931
- 廣東 (Caradja & Meyrick, 1934); 廣州 (Shibuya, 1931).
- Sy. aurantiacalis* Fisch v. Rösl.
- 中國 (Hampson, 1896); 華中、華西、福建、台灣、廣州 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja, & Meyrick, 1934).
- Sy. balteata* (Fabricius, 1798)
- 華中、華南、華西 (Klima, 1939); 浙江 (李鳳藻, 1952) 台灣 (Shibuya, 1928; Klima, 1939).
- Sy. bicolor* Caradja

- 廣東、四川 (Caradja & Meyrick, 1934).
- Sy. cantonalis* Shibuya, 1931
廣州 (Shibuya, 1931).
- Sy. capnosalis* Caradja, 1925
華南、華西 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Sy. chalybifascia* Hampson, 1896
台灣等 (Shibuya, 1928; Klima, 1939).
- Sy. cometa tamsi* Caradja, 1927
中國 (Caradja, 1927); 華西 (Klima, 1939).
- Sy. concatenalis* Walker
廣東 (Caradja & Meyrick, 1934).
- Sy. costalis* (Moore, 1888)
華南 (Klima, 1939); 華西 (Caradja, 1925; Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939);
廣東 (Caradja 1925; Caradja & Meyrick, 1934);
- Sy. crotonalis* (Walker, 1859)
台灣等 (Shibuya, 1928; Klima, 1939).
- Sy. deficiens* (Moore, 1887)
華南 (Klima, 1939); 華東、華西 (Caradja, 1925; Klima, 1939); 台灣 (Shibuya, 1928; Klima,
1934); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Sy. delicatalis* Strand, 1918
台灣 (Klima, 1939).
- Sy. derogata* (Fabricius, 1775)
中國 (Hampson, 1896 (*multilinealis*)); 華北、華南 (Klima, 1939); 華中、華西 (Caradja, 1925;
Klima, 1939); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1939; 李鳳藻, 1952); 廣東 (Caradja &
Meyrick, 1934); 上海、江西、廣州 (Caradja, 1925); 河北、山西、陝西、山東、江蘇、浙江、安徽、
江西、湖南、湖北、河南、福建、廣西、雲南、四川、貴州 (李鳳藻, 1952).
- Sy. dissipatalis* (Lederer, 1863)
華南、台灣、(Klima, 1939); 福建、廣州 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja &
Meyrick, 1934).
- Sy. distinguenda* Hering
廣東等 (Caradja & Meyrick, 1934).
- Sy. euryterminalis* Hampson, 1918
華南 (Klima, 1939); 台灣 (Shibuya, 1928; Caradja & Meyrick, 1934; Klima, 1939); 廣東
(Caradja & Meyrick, 1934).
- Sy. fabiusalis makanalis* Caradja, 1932
華南 (Caradja, 1932); 浙江 (Klima, 1934).
- Sy. fraterna* Moore
廣東等 (Caradja & Meyrick, 1934).
- Sy. fuscomarginalis* (Leech, 1889)
華中、華南 (Klima, 1939); 華西 (Caradja, 1925; Klima, 1925); 浙江、福建 (Caradja, 1925);
廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Sy. haryoalis Strand, 1918

台灣 (Shibuya, 1928; Klima 1939).

Sy. hoenei Caradja, 1925

福建 (Caradja, 1925; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934; Klima, 1939).

Sy. imbutalis taihoqualis Strand, 1918

台灣 (Shibuya, 1928; Klima, 1939).

Sy. inferior Hampson, 1898

華中 (Caradja, 1925; Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939).

Sy. insignis (Butler, 1881)

華中、華西 (Caradja, 1925; Klima, 1939).

Sy. invalidalis South, 1901

中國 (South, 1901); 華中、華西、江西 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Sy. topasalis (Walker, 1859)

華中、華西 (Caradja, 1925; Klima, 1939); 華南 (Klima, 1939); 台灣 (Hampson, 1896; Caradja, 1925; Shibuya, 1928; Klima, 1939); 福建、廣州 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Sy. topasalis f. plagialis Walker, 1859

廣東 (Caradja & Meyrick, 1934).

Sy. topasalis leopardalis Moore, 1888

台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).

Sy. karenkonis Shibuya, 1928

台灣 (Shibuya, 1928; Klima, 1939).

Sy. lucidalis Caradja, 1925

華中、華西 (Klima, 1939); 上海 (Caradja, 1925).

Sy. luctuosalis (Guenée 1854)

中國 (Hampson, 1896); 華中 (Caradja, 1925; Klima, 1939); 華南 (Klima, 1939); 華西、上海 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 江蘇、浙江 (李鳳蓀, 1952).

Sy. lulalis Strand, 1918

台灣 (Shibuya, 1928; Klima, 1939).

Sy. lunalis (Guenée, 1854)

台灣等 (Hampson, 1896; Shibuya, 1929; Klima, 1939).

Sy. lutealis South

華中 (Caradja, 1925).

Sy. luteolalis South, 1901

中國 (South, 1900); 華西 (Klima, 1939).

Sy. maculalis (Leech, 1889)

華中、華西 (Caradja, 1925; Klima, 1939); 華南 (Klima, 1939); 福建、台灣 (Caradja 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Sy. magna (Butler, 1879)

華中 (Caradja, 1925; Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Sy. mandarinialis Caradja, 1925

中國 (Caradja, 1925; Klima, 1939).

Sy. moderatalis (Christoph, 1881)

華中、華南、華西 (Klima, 1939).

Sy. mundalis (South, 1901)

台灣等 (Shibuya, 1928; Klima, 1939).

Sy. mysialis (Walker, 1859)

華中、華南、華西 (Klima, 1939); 上海、福建、廣東、廣州 (Caradja, 1925).

Sy. mysialis var. *impuralis* (Warren, 1896)

廣東 (Caradja & Meyrick, 1934).

Sy. ningpoalis (Leech, 1889)

華中 (Caradja, 1925; Caradja & Meyrick, 1934; Klima, 1939); 華東 (Caradja, 1925; Klima, 1939); 華南 (Caradja & Meyrick, 1934; Klima, 1939); 華西 (Klima, 1939); 浙江、福建、廣州 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Sy. orobentalis Snellen

廣州 (Caradja, 1925).

Sy. paucistrialis (Warren, 1896)

華中、華南 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Sy. pernitescens (Swinhoe, 1894)

華南 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 廣州 (Caradja 1925); 台灣 (李鳳蓀 1952).

Sy. proctizonalis Hampson, 1918

華中 (Klima, 1939).

Sy. pseudovalis Hampson

廣東 (Caradja & Meyrick, 1934).

Sy. quadrimaculalis (Kollar, 1844)

華南 (Klima, 1939); 華中、華西 (Caradja, 1925; Klima, 1939); 江西、福建、廣州 (Caradja, 1925); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).

Sy. rhyparialis Oberthür, 1893

華中、華西 (Caradja, 1925; Klima, 1939).

Sy. ruralis (Scopoli, 1763)

華西 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939).

Sy. ruralis var. *dubia* (Hampson, 1891)

廣東 (Caradja & Meyrick 1934).

Sy. ruricolalis Snellen

廣東 (Caradja & Meyrick, 1934).

Sy. sabinusalis (Walker, 1859)

- 華中、華西、台灣 (Caradja, 1925; Klima, 1939).
- Sy. signalis* (Leech, 1889)
華中等 (Caradja, 1925; Klima, 1939).
- Sy. sellalis* (Guenée, 1854)
華西、台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).
- Sy. sublittoralis* (Walker, 1865)
華南等 (Klima, 1939).
- Sy. taiwanalis* Shibuya, 1928
台灣等 (Shibuya, 1928; Klima, 1939).
- Sy. tricolor* (Butler, 1879)
華中、華西 (Caradja, 1925; Klima, 1939); 台灣 (Shibuya, 1928); 廣東、四川 (Caradja & Meyrick, 1934).
- Sy. ultimalis* (Walker, 1859)
華南 (Klima, 1939); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934 (var.)); 廣州 (Caradja, 1925 (var.)).
- Sy. venustalis* Swinhoe, 1894
華北、華南、華西 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Syngamia abruptalis* (Walker, 1859)
台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).
- Syn. falsidicalis* (Walker, 1859)
華中 (Caradja, 1925; Klima, 1939); 華南、華西 (Klima, 1939); 江西、廣州 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).
- Syn. floralis* (Zeller, 1852)
華南 (Klima, 1939); 台灣 (Caradja, 1925; Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Syn. latimarginalis* (Walker, 1859)
台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).
- Syn. vibiusalis* (Walker, 1859)
台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja & Meyrick, 1934).
- Tabidia candidalis* (Warren, 1896)
華南 (Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- T. strigiferalis* Hampson, 1900
華南、華西 (Klima, 1939); 福建、廣州 (Caradja, 1925); 廣東 (Caradja & Meyrick, 1934).
- Terastia meticulosalis* Guenée, 1854
台灣等 (Shibuya, 1928).
- Te. subjectalis* Lederer
廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925 (*Terastia*)).
- Te. vinacealis* (Moore, 1877)
福建 (Caradja, 1925 (*calceolaris*)); 台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).
- Thliptoceras octoguttale* Felder
廣東 (Caradja & Meyrick, 1934).

- Th. stygale* Hampson, 1896
廣東 (Caradja, 1925).
- Th. variabilis* Swinhoe, 1890
福建、廣州 (Caradja, 1925(*variabile*)); 台灣 (Shibuya, 1928); 廣東 (Caradja & Meyrick, 1934).
- Titanio basalis* Caradja, 1928
新疆 (Caradja, 1928).
- Tyspanodes hypsalis* Warren, 1891
華北 (Klima, 1939); 華西 (Caradja, 1925; Klima, 1939); 上海、南京 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima 1939).
- Ty. linealis* Moore 1867
華南 (Klima, 1939); 廣東 (Caradja & Meyrick, 1934); 廣州 (Caradja, 1925).
- Ty. obscuralis* Caradja, 1925
華東 (Caradja & Meyrick, 1934); 華南 (Caradja & Meyrick, 1934; Klima, 1939); 廣州 (Caradja, 1925); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Ty. striata* (Butler, 1879)
華南、華西 (Klima, 1939); 江西 (Caradja, 1925); 台灣 (Shibuya, 1928; Klima, 1939); 廣東 (Caradja, 1925; Caradja & Meyrick, 1934).
- Ty. striata* var. *saturata* Caradja, 1925
華中、華南 (Klima, 1939); 南京 (Caradja, 1925).
- Ulopeza flavomaculata* Shibuya, 1931
廣州 (Shibuya, 1931).
- U. idyalis* (Walker, 1859)
華南 (Klima, 1939); 福建 (Caradja, 1925).
- U. semivialis nigrocotalis* Caradja, 1925
廣東 (Caradja, 1925; Caradja & Meyrick, 1934); 華南 (Klima, 1939).

參 考 文 獻

- [1] 李鳳藻, 1952, 中國經濟昆蟲學, (中卷).
- [2] 錢念曾, 1940 (?), 我國之倉庫害蟲, 湖南農業改進所.
- [3] Butler, A. G., 1888, On the Lepidoptera recieved from Dr. Emin Pasha. *Proc. Zool. Soc. Lond.* 1888: 56-85.
- [4] Caradja, A., 1925. Ueber Chinas Pyraliden, Tortriciden, Tineiden, nebst kurze Betrachtungen, zu deren das Studium dieser Fauna Verablassung gibt. *Acad. Romana, Mem. Sect. stint. ser.* iii, 3(7): 257-383.
- [5] ———, 1926. Ueber Chinas Pyraliden, Tortriciden, Tineiden, nebst kurze Betrachtungen, zu deren das Studium dieser Fauna Veranlassung gibt. *Bull. Acad. Roumaine*, 10(2): 3-6.
- [6] ———, 1927. Die Kleinfalter der Stötzner'schen Ausbeute, nebst Zuträge aus meiner Sammlung. *Acad. rom., Mem. Sect. stint.*, 3(4), mem. 8, 68 pp.
- [7] ———, 1931. Second contribution to our knowledge about the Pyralidae and Microlepidoptera of Kwanhsien. *Bull. Acad. Romaine*, 14: 59-75.
- [8] ———, 1932. Beiträge zur Lepidoptera-Fauna Grossrumäniens für das Jahr 1931. *Bull.*

- Acad. Romnatne*, 15: 35-46.
- [9] ———, 1938. Materialien zu einer Microlepidoptera-Fauna von Atuntse in Nord-Yunnan. Ein Nachtrag zur gleichbetitelten arbeit in D. E. Z. *Iris Stettin ent. Ztg.* 99: 247-250.
- [10] ———, 1938. Beiträge zur Lkeinfalterfauna Chinas (Lep.). *Ibid.*, 99: 250-253.
- [11] ———, 1938. Materialien zu einer Microlepidopteren-Fauna Nord-Fukiens. *Ibid.* 99: 253-257.
- [12] ———, 1939. Materialien zu einer Mikrolepidopterenfauna von Atuntse (Nordwest-Yünnan). *Iris* 52:99-104.
- [13] ———, 1939. Materialien zu einer Lepidopterenfauna des Taipeishanmassivs (Tsiulinshan), Provinz Shensi. *Iris* 52: 104-111.
- [14] ———, 1939. Materialien zu einer Mikrolepidopteranfauna des Mienshan, Provinz Shansi, China. *Iris* 53: 1-15.
- [15] ———, 1939. Materialien zu einer Mikrolepidopterenfauna des Yangtsetales bei Batang. *Iris* 53: 15-26.
- [16] ———, 1939. Materialien zu einer Mikrolepidopterenfauna von Kuatun in der chinesischen Provinz Fukien. *Iris* 53: 27-32.
- [17] Caradja, A. und E. Meyrick, 1933. Materialien zu einer Microlepidopteren Fauna Kwangtungs. *Iris* 47: 123-144.
- [18] ———, 1936. Materialien zu einer Lepidopterenfauna des Taishanmassivs, Provinz Shangtung. *Iris* 50: 35-46.
- [19] ———, 1936. Materialien zu einer Lepidopterenfauna des Taishanmassivs, Provinz Shangtung. (Contd.) *Iris* 50: 145-59.
- [20] ———, 1937. Materialien zu einer Mikrolepidopteren Fauna des Yüling-shanmassivs (Provinz Yünnan). *Iris* 51: 137-82.
- [21] ———, 1938. Materialien zu einer Mikrolepidopterafauna des Yüling-shanmassivs (Provinz Yünnan). *Iris* 52: 1-29.
- [22] Hampson, G. F. 1895. On the classification of the Schoenobiinae and Crambinae, two subfamilies of moths of the family Pyralidae. *Proc. Zool. Soc. Lond.* 1895: 897-974.
- [23] ———, 1896, On the classification of three subfamilies of moths of the family Pyralidae: the Epipaschiinae, Endotrichinae and Pyralinae. *Trans. ent. Soc. Lond.* 1896: 451-550.
- [24] ———, 1897, On the classification of the two subfamilies of moths of family Pyralidae: the Hydrocampinae and Scopariinae. *Trans. ent. Soc. Lond.* 1897: 127-240.
- [25] ———, 1899. The moths of India. Supplementary paper to the Volumes in "the fauna of British India" Part VI. *J. Bombay Soc.* 12: 475-85; Part VII, tc. pp. 697-715.
- [26] ———, 1898. A revision of the moths of the subfamily Pyraustinae and family Pyralidae Part I. *Proc. Zool. Soc. Lond.* 1898: 590-761.
- [27] ———, 1899. A revision of the moths of the subfamily Pyraustinae and family Pyralidae Part II. *Ibid.* 1899: 172-291.
- [28] Kato, Shizuo, 1949, A preliminary report on a survey of agricultural insect pests in Chahar, Suiyuan and Northern Shanshi. *Pek. nat. Hist. Bull.* 18 (1): 11-36.
- [29] Klima, A., 1937, Pyralididae: Subfamily: Scopariinae, Nymphulinae in Bryk's Lepidop-

terorum Catalogus 84, 226 pp.

- [30] ———, 1939, Pyralididae: Subfamily: Pyraustinae I in Bryk's *Lepidopterorum Catalogus* 89, 224 pp.
- [31] Leech, J. H. *Lepidoptera Heterocera from China, Japan and Corea. Part V. Trans. ent. Soc. Lond.* 1901: 385-514.
- [32] Meyrick, E. 1937. *Exotic Microlepidoptera* 5: 65-160.
- [33] Shibuya, J. 1928. The systematic study of the Formosan Pyralidae. *J. Fac. Agric. Sapporo* 22: 1-300.
- [34] Strand, R., 1919, Sauter's Formosa-Ausbeute: Pyralididae, Subfamily Galleriinae, Crambinae, Schoenobiinae, Anerastiinae und Phycitinae. *Ent. Zeitung Stettin* 79: 248-76.
- [35] ———, 1922, Sauter's Formosa-Ausbeute. *Ent. Zeitschr. Frankfurt*. 36: 1-2, 6-7, 11, 15, 19.
- [36] West, R. J., 1931, Descriptions of new species of Japanese, Formosan and Philippine Pyralidae. *Novit. Zool.* 36: 206-19.
- [37] Wileman, A. E. and R. South, 1919. New species of Pyralidae from Formosa. *Ent. Lond.* 52: 267-68.
- [38] Wu, C. F., 1938. Pyralidae in *Catalogus Insectorum Sienesium*, Vol. IV, pp. 63-132.